

# Real Number System

## (Equality and Inequality Properties)

**Our current goal is to list the equality and inequality properties the real numbers possess.**

We have encountered various properties the set of real numbers possesses as we considered its type, operations, ... Now, let's list and name these properties that involve equality ( = ) in one place.

Assume that  $a$ ,  $b$ , and  $c$  are real numbers:

### Equality Properties:

Real Number Properties	Comments	Examples
<b>1. Closure</b>		
a. $a + b$ is a real number	The <i>sum</i> of two real numbers is a real number	$2 + 5 = 7$ is a real number
b. $a \cdot b$ ( = $a b$ ) is a real number	The <i>product</i> of two real numbers is a real number	$2 \cdot 5 = 10$ is a real number
<b>2. Commutative Property</b>		

a. $a + b = b + a$	Two real numbers can be <i>added</i> in either order	$2 + 3 = 5 = 3 + 2$
b. $a * b = b * a$	Two real numbers can be <i>multiplied</i> in either order	$2 * 7 = 14 = 7 * 2$
	Note: $a - b$ is probably not $b - a$	$5 - 2 = 3$ $2 - 5 = -3$
<b>3. Associative Property</b>		
a. $(a + b) + c = a + (b + c)$	The <i>sum</i> of three real numbers does not depend on the order they are <i>added</i> .	$(4 + 3) + 5 = 7 + 5 = 12$ $4 + (3 + 5) = 4 + 8 = 12$
b. $a *(b*c) = (a*b)*c$	The <i>product</i> of three real numbers does not depend on the order they are <i>multiplied</i> .	$2*(4*3) = 2*12 = 24$ $(2*4)*3 = 8*3 = 24$
<b>4. Identity Property</b>		
a. $a + 0 = a = 0 + a$	"0" is called the <b><i>additive identity</i></b> . The <i>sum</i> of a real number and "0" is the number itself.	$6 + 0 = 6 = 0 + 6$
b. $a * 1 = a = 1 * a$	"1" is called the <b><i>multiplicative identity</i></b> . The <i>product</i> of a real number and "1" is the number itself	$9 * 1 = 9 = 1 * 9$
<b>5. Inverse Properties</b>		

<p>a. For each real number “a”, there is a real number “- a” such that</p> $a + (- a) = 0$ $= (- a) + a$	<p>“- a” is called the <b>additive inverse</b> of “a”. The <i>sum</i> of a real number and its additive inverse is “0”</p> <p>Note: a + (- a) is frequently written a - a</p>	$4 + (-4) = 0 = (-4) + 4$
<p>b. For each real number a (not equal to 0), there is a real number “1/a” such that</p> $a*(1/a) = 1$ $= (1/a)*a$	<p>(1/a) is called the <b>multiplicative inverse</b> of “a”. The <i>product</i> of a real number and its multiplicative inverse is “1”</p>	$8*(1/8) = 1 = (1/8)*8$
<p><b>6. Distributive Property</b></p>		
<p>a. <math>a*(b + c) =</math> <math>a*b + a*c</math></p>	<p>The <i>product</i> of a real number and the <i>sum</i> of two real numbers is the <i>sum</i> of the <i>product</i> of the first number and each of the other numbers.</p>	$3*(4 + 2) = 3*6 = 18$ $3*4 + 3*2 = 12 + 6 = 18$
<p>b. <math>a*(b - c) =</math> <math>a*b - a*c</math></p>	<p>The <i>product</i> of a real number and the <i>difference</i> of two real numbers is the <i>difference</i></p>	$3*(4 - 2) = 3*2 = 6$ $3*4 - 3*2 = 12 - 6 = 6$

	of the <i>product</i> of the first number and each of the other numbers.	
<b>7. Multiplicative Property of Zero</b>		
$a \cdot 0 = 0 \cdot a = 0$	The <i>product</i> of a real number and "0" is "0"	$2 \cdot 0 = 0 = 0 \cdot 2$

**Inequality Properties:**

Terms (“+” or “-”)		
<p style="text-align: center;"><b>Addition</b></p>	<p>1. If <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p style="text-align: center;"><math>a + c \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b + c</math></p>	<p><math>6 \leq 8</math> implies <math>6 + 3 = 9 \leq 11 = 8 + 3</math></p>
<p style="text-align: center;"><b>Subtraction</b></p>	<p>2. If <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p style="text-align: center;"><math>a - c \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b - c</math></p>	<p><math>6 \leq 8</math> implies <math>6 - 3 = 3 \leq 5 = 8 - 3</math></p>

Factors ("*" or "/")		
	<p>1. If <math>c &gt; 0</math> then</p> <p>a. <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p><math>a * c \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b * c</math></p>	<p><math>3 \leq 7</math></p> <p>implies</p> <p><math>3 * 2 = 6 \leq 14 = 7 * 2</math></p>
	<p>b. <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p><math>\frac{a}{c} \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} \frac{b}{c}</math></p>	<p><math>3 \leq 7</math></p> <p>implies</p> <p><math>\frac{3}{2} \leq \frac{7}{2}</math></p>

	2. If $c < 0$ then	
	<p>a. <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p><math>a * c \left\{ \begin{array}{l} \geq \\ &gt; \\ \leq \\ &lt; \end{array} \right\} b * c</math></p>	$3 \leq 8$ implies $3 * (-2) = -6 \geq -16 = 8 * (-2)$
	<p>b. <math>a \left\{ \begin{array}{l} \leq \\ &lt; \\ \geq \\ &gt; \end{array} \right\} b</math> then</p> <p><math>\frac{a}{c} \left\{ \begin{array}{l} \geq \\ &gt; \\ \leq \\ &lt; \end{array} \right\} \frac{b}{c}</math></p>	$3 \leq 8$ implies $3 / (-2) = -\frac{3}{2} \geq -4 = 8 / (-2)$

**WARNING:** When multiplying or dividing by a negative real number, the direction (“sense”) of the inequality switches; for example “ $\leq$ ” changes to “ $\geq$ ”. **PLEASE Be Careful!**